

## **Edexcel A-Level Maths 2026 Paper 2**

Do not turn over the page until instructed to do so.

This assessment is out of 100 marks and you will be given 120 minutes.

This is **not a predicted paper** but is an opportunity to practice some topics likely to come up.


When you are asked to by your teacher write your **full name** below

**Name:**

**Total Marks:**                      **/ 100**

**1**

The figure below shows a sketch of a curve  $C$  and a straight line  $l$ .



Given that

- $C$  has an equation  $y = f(x)$  where  $f(x)$  is a quadratic expression in  $x$ .
- $C$  cuts the  $x$ -axis at 1 and 5.
- $l$  cuts the  $x$ -axis at  $(1,0)$  and intersects the curve at the point  $(4,6)$ .

Use inequalities to define the region  $r$ .

**(5)**



**2**

Given that  $\theta$  is measured in radians, prove, from first principles, that the derivative of  $\sin(\theta)$  is  $\cos(\theta)$ .

You may assume that as  $h \rightarrow 0$ ,  $\frac{\sin(h)}{h} \rightarrow 1$  and that  $\frac{\cos(h) - 1}{h} \rightarrow 0$

**(5)**

**3**

- a)** Given that both  $(x + 1)$  and  $(x - 3)$  are both factors of

$$p(x) = x^4 + 5x^3 + ax^2 + bx - 36,$$

find the values of  $a$  and  $b$ .

**(4)**

- b)** Hence, fully factorise  $p(x)$  as a product of linear factors.

**(3)**

- c)** Sketch  $p(x)$ .

**(2)**



**4**

Jess is modelling the height,  $h$ , of a daffodil at the time  $t$ , in weeks, after the shoot appeared.

She models it with the function

$$h^2 = at + b, \quad 0 \leq t \leq 6$$

Given that

- the height of the daffodil was 15 cm at  $t = 2$ .
- the height of the daffodil was 18 cm at  $t = 3$ .

**a)** Find a complete equation for the model, giving the values of  $a$  and  $b$  to three significant figures.

**(4)**

**(b)** Evaluate this model's suitability as a model for the whole of the time period.

**(2)**



**5** In a secluded woodland some rare fungus grows on an oak stump.

The area,  $A$  cm<sup>2</sup>, of the fungus is measured for a few days and is shown to follow the model

$$A = 3 + 5e^{0.2t}$$

where  $t$  is the number of days after the fungus was first observed.

**a)** Sketch a graph of  $A$  against  $t$ , showing any intersections with the coordinate axes.

**[1 mark]**

**b)** Find the number of days for the area of the fungus to triple in size.

**[3 marks]**

**c)** Critique the model.

**[1 mark]**



**6** Solve,

**a)** Solve  $2 \log_2(x + 3) - \log_2(2x) = 3$

**(2)**

**b)** Use the substitution  $y = 2^{3x}$  to solve


$$64^x - 2^{3x+2} - 5 = 0$$

**(3)**



7

The graph of  $y = \frac{20}{1 + \sqrt{x}}$  is shown below.



- a) Use the Trapezium Rule with 7 ordinates to find an approximation to the shaded area.

(4)

- b) Use your answer to part (a) to state an approximate value of  $\int_2^8 \frac{10}{1 + \sqrt{x}} dx$

(1)

- c) By finding analytically  $\int_2^8 \frac{10}{1 + \sqrt{x}} dx$ , find the percentage error made in the approximation performed in parts (a) and (b).

(7)

(d) Use part (a) to find  $\int_6^{12} \frac{20}{1 + \sqrt{x+4}}$

(2)



**8**

Geoff bought a classic car in 1996 for £9800.

A classic car specialist valued the car at 5 year intervals as shown in the table below.


<b>Year</b>	1996	2001	2006	2011
<b>Value</b>	9800	19500	39100	79000

The valuer suggests that the valuation price can be modelled by the equation  $V = a \times b^t$ , where  $t$  is the number of years after 1996.

- a)** Find the linearised form of the model given above. **(2)**
- b)** Complete the table below:

$t$	0	5	10	15
$\log_{10}(V)$				

- (2)**
- c)** By plotting a graph on the axes below, estimate the values of  $a$  and  $b$ . **(4)**
- d)** Find the expected value of the car in 2032 and comment on the validity of this value. **(2)**





**9**Find the points of inflection of the curve with equation  $y = \cot(2x)$ **(7)**



10

a) Find, in ascending powers of  $x$ , up to the term including  $x^3$ , the binomial expansion of  $(1 + x)^{-\frac{1}{3}}$

(2)

b) A student approximates  $\frac{1}{\sqrt[3]{7}}$  using the expansion found in (a) and the value  $x = 6$ . Explain what is wrong in doing this.

(1)

c) By using a suitable value of  $x$  and part (a), find an approximation to  $\frac{1}{\sqrt[3]{7}}$ .

(4)



**11**

The rate of change of the radius of a circle is inversely proportional to the radius cubed.

- a)** Show that the rate at which the area of the circle,  $A$ , changes

satisfies 
$$\frac{dA}{dt} = \frac{2\pi^2}{A}$$

**(4)**

- b)** Explain why  $\frac{dA}{dt} > 0$

**(1)**



**12**

Consider the circle  $x^2 - 8x + y^2 - 8y + 7 = 0$

**a)** What is the centre,  $C$ , of the circle?

**[1 mark]**

**b)** Find the equation of the tangent to the circle at  $P(8,7)$

**[3 marks]**

**c)** Another tangent to the circle at  $Q(7,0)$  has equation  $3x - 4y = 21$ .

This tangent meets the tangent found in (a) at the point  $R$ .

Shape  $T$  is formed by removing the sector  $CPQ$  of the circle from the quadrilateral  $CPRQ$ .

Find the ratio

Perimeter of  $T$  : Area of  $T$


**[8 marks]**





**13**

The sketch below shows the curve with equation  $x^2 + xy + y^2 = a$  for some constant  $a$ .



Find the area of the rectangular box that just contains the curve.

**(10)**

